

Syntheses and Characterizations of Some New N-alkyl, Isoxazole and Dioxazole Derivatives of 5-Chloroisatin

Z. Tribak¹, M.K. Skalli^{1*}, O. Senhaji², Y. Kandri. Rodi¹

¹Laboratory of Applied Chemistry, Sidi Mohamed Ben Abdellah University, Faculty of Sciences and Technology of Fez, Morocco.

²Laboratory of Applied Physical Chemistry, Moulay Ismaïl University, Faculty of Sciences and Technology of Errachidia, Morocco.

Abstract— *N-alkyl and cycloadducts derivatives of 5-Chloroisatin were synthesized in good to excellent yields. The method evidences a selective N-alkylation when using 1,2-bis (2-chloroethoxy) ethane as efficient spacer at room temperature on the 5-Chloroisatin moiety. A general method for the 1,3-dipolar cycloaddition of 4-Chlorobenzaldoxime to alkynes provides a useful alternative route to get new isoxazole et dioxazole derivatives.*

Keywords— *1, 3-dipolar cycloaddition, 4-Chlorobenzaldoxime, N-alkylation, ¹H NMR, ¹³C NMR.*

I. INTRODUCTION

The interest for the synthesis of the isatin derivatives has increased in organic and pharmaceutical chemistry. In 1988 the isatin is isolated and reported to possess a wide range of activities involving the central nervous system as an endogenous compound [1-3]. This heterocyclic system is found in a number of molecules possessing biological and pharmacological properties. The usage of some isatin derivatives for antibacterial, antimalarial, anti-proliferative and anti-coagulant [4-9], antimicrobial, anticonvulsant, anticancer, antiHIV, etc. activities and also antifungal, herbicidal agents [10-17]. Many other remarkable applications are reported in the literature, such as the selective inhibition of corrosion in mild steel by 5-chloro-1-(2-(dimethylamino)ethyl)indoline-2,3-dione and 5-chloro-1-octylindoline-2,3-dione [18, 19]. These 5-Chloroisatin derivatives were synthesized using N-alkylation reaction under catalysis by phase transfer [20-23].

Tribak et al. have been prepared new 5-Chloroisatin derivatives such as 5-chloro-1-((3-(4-chlorophenyl)-4,5-dihydroisoxazol-5-yl) methyl) indoline-2,3-dione and 5-chloro-3'-(4-chlorophenyl)-1-((3-(4-chlorophenyl)-4,5-dihydroisoxazol-5-yl) methyl) spiro [indoline-3,5'-[1,4,2]dioxazol]-2-one that would carry medicinal properties of mainly as antibacterial against two bacteria Gram+ and

two other bacteria Gram- [24, 25]. These derivatives have also been synthesized by 1,3-dipolar cycloaddition between nitrile oxides and N-alkylchloroisatin using Sodium hypochlorite [26]. Therefore, as part of our ongoing study, we report herein the synthesis, and characterization of new N-alkylchloroisatin by N-alkylation method and also new isoxazole and dioxazole compounds formed from the condensation of 4-Chlorobenzaldoxime with 5-chloro-1-(prop-2-yn-1-yl) indoline-2,3-dione.

II. EXPERIMENTAL DETAILS

All chemicals were purchased from sigma - Aldrich and used as such. Solvents hexane, methanol and CHCl₃ (AR grade) were used without further purification.

The reactions were monitored by thin-layer chromatography (TLC) analysis by using silica gel (60 F254) plates. Compounds were visualized by UV irradiation at 256 or 365 nm. Flash column chromatography was performed on silica gel 60 (230–400 mesh, 0.040–0.063 mm). Melting points (m.p. [°C]) were taken on samples in open capillary tubes and are not corrected

¹H and ¹³C NMR spectra were recorded with a Bruker spectrometer at 300 and 75 MHz respectively. Chemical shifts are given in parts per million (ppm) from tetramethylsilane (TMS) as internal standard in CDCl₃, and the residual peak of DMSO in [D₆] DMSO. The following abbreviations are used for the ¹H NMR spectra multiplicities: br. s: broad singlet, s: singlet, d: doublet, t: triplet, q: quartet, qt: quintuplet, m: multiplet. Coupling constants (J) are reported in Hertz [Hz].

2.1. Preparation of the compound (1a):

To a solution of (0.4 g, 2.20 mmol) 5-chloro-1H-indole-2,3-dione in 20 ml of DMF, 0.5 g, 3.3 mmol of K₂CO₃ and 0.1 g, 0.3 mmol of tetra-n-butyl ammonium bromide (BTBA), 0.20 mL (1.32 mmol) of 1,2-bis (2-chloroethoxy) ethane is added dropwise.


The reaction was refluxed with DMF for 48 hours. After removal of the salts by filtration, the DMF is evaporated under reduced pressure and the residue obtained is dissolved in dichloromethane, the mixture obtained is chromatographed on a silica gel column (eluent: ethyl acetate / Hexane (4/1)). The reaction allowed the solid 22 to be isolated

1-(2-(2-(2-chloroethoxy)ethoxy)ethyl)-5-chloroindoline-2,3-dione (1a): Yield: 88% ; mp: 90-120; $R_f = 0.85$; (eluent Ethyl acetate / hexane (4: 1)). $^1\text{H NMR}$ (CDCl_3 ; 300MHz) : δ (ppm) 7.49-7.52 (m, 2H, H_{Ar}); 7.05 (d, H, H_{Ar} , $^3J_{\text{H-H}} = 6\text{Hz}$,); 3.91 (t, 2H, CH_2 , $^3J_{\text{H-H}} = 6\text{Hz}$, $^4J_{\text{H-H}} = 3\text{Hz}$); 3.75 (t, 2H, CH_2 , $^3J_{\text{H-H}} = 6\text{Hz}$, $^4J_{\text{H-H}} = 3\text{Hz}$); 3.67 (t, 2H, CH_2 , $^3J_{\text{H-H}} = 6\text{Hz}$, $^4J_{\text{H-H}} = 3\text{Hz}$); 3.59 (m, 4H, CH_2); 3.21 (t, 3H, CH_3 , $^3J_{\text{H-H}} = 6\text{Hz}$, $^4J_{\text{H-H}} = 3\text{Hz}$). $^{13}\text{C NMR}$ (CDCl_3 ; 75MHz): δ (ppm) 183.11 (C=O); 167.00 (N-C=O); 143.67, 130.74, 119.51 (Cq); 134.98, 130.32, 123.75 (CH_{Ar}); 71.38, 63.11, 40.85, 20.92 (CH_2).

2.2. General procedures for cycloaddition of N-propargylchloroisatin with nitrile oxide:


0.2 g of 5-chloro-1-(prop-2-yn-1-yl)indoline-2,3-dione and (0.17 g, 1.079 mmol) of 4-Chlorobenzaldoxime were dissolved in 12 mL of chloroform (CHCl_3) when the mixture reaches 0°C , 4 mL of bleach (NaOCl) was introduced and then allowed to stir for 4 hours. The resulting compound purified and recrystallized from ethanol.

5-chloro-1-((3-(4-chlorophenyl)isoxazol-5-yl)methyl)indoline-2,3-dione (1b): Yield: 77%; mp:



Scheme 1

The 1,3-dipolar cycloaddition reaction of nitrile oxide, prepared in situ by the action of bleach on 4-Chlorobenzaldoxime, with 5-chloro-1-(prop-2-yn-1-yl)indoline-2,3-dione in a biphasic medium



Scheme 2

194-197 ; $R_f = 0.77$; (eluent Ethyl acetate / hexane (6: 1)). $^1\text{H NMR}$ (CDCl_3) δ ppm 7.71-7.81 (m, 2H, H_{Ar}); 7.57 (d, H, H_{Ar} , $J = 8.4\text{Hz}$); 6.97-7.04 (m, 4H, H_{Ar}); 6.55 (s, H, CH); 3.87 (m, H, CH_2). $^{13}\text{C NMR}$ (CDCl_3) δ ppm: 185.86 (C=O) 160.64 (N-C=O); 163.18, 157.88, 147.07, 134.14, 131.17, 130.95, 118.02 (Cq) 134.56, 130.11, 129.26, 128.62, 123.96 (CH_{Ar}); 105.51 (CH); 50.60 (CH_2)

5-chloro-3'-(4-chlorophenyl)-1-((3-(4-chlorophenyl)isoxazol-5-yl)methyl)spiro[indoline-3,5'[1,4,2]dioxazol]-2-one (2b): Yield: 65%, mp: 197-199; $R_f = 0.75$; (eluent Ethyl acetate /hexane (5: 1)). $^1\text{H NMR}$ (CDCl_3) δ ppm 7.80 (d, 2H, H_{Ar} , $J = 9\text{Hz}$); 7.73 (d, 2H, H_{Ar} , $J = 8.4\text{Hz}$); 7.56 (d, H, H_{Ar} , $J = 1.8\text{Hz}$); 7.48-7.46 (m, 5H, H_{Ar}); 7.05 (d, H, H_{Ar} , $J = 8.4\text{Hz}$); 6.59 (s, H, CH); 5.05 (s, 2H, CH_2). $^{13}\text{C NMR}$ (CDCl_3) δ ppm: 168.12 (NC=O); 166.03, 162 (C=N); 141.45, 138.63, 136.51, 130.20, 126.77, 122.62, 120.05 (Cq); 133.39, 128.52, 128.10, 126.65, 111.11 (CH_{Ar}); 101.60 (CH); 35.71 (CH_2).

III. RESULTS AND DISCUSSION

Based on the interest of our research group in the synthesis of new heterocycles [27], we studied the condensation of 5-chloroisatin with 1,2-bis(2-chloroethoxy) ethane under the conditions of liquid/solid phase transfer catalysis in DMF as solvent and potassium bicarbonate as a weak base at 80°C allowed us to isolate compound 5-chloro-1-(2-(2-(2-chloroethoxy)ethoxy)ethyl)indoline-2,3-dione (scheme 1)

(water/ethanol) at 0°C for 4 hours. It leads in each case to the formation of the two cycloadducts resulting from the 1,3-dipolar cycloaddition on the dipolarophile group C-O in position 3 and the propargyl group in position 1.

IV. CONCLUSION

In conclusion, we have developed an efficient synthetic method for the selective N-alkylation of 5-Chloroisatin using an easily available mild base K_2CO_3 by the action of 1,2-bis (2-chloroethoxy) ethane. Subsequently, an easy and general 1,3-dipolar cycloaddition method has also been demonstrated using nitric oxide at 0°C temperature with 5-chloro-1-(prop-2-yn-1-yl)indoline-2,3-dione as dipolarophile allows getting new cycloadducts containing the two nuclei isoxazole and dioxazole. Their structures has been identified by the usual spectroscopy.

ACKNOWLEDGEMENTS

The authors would like to thank all the people who helped to carry out this work such as 1H NMR, ^{13}C NMR.

REFERENCES

- [1] Tribak, Z.; Skalli, M. K.; Senhaji, O.; KandriRodi, Y.; DFT Study and Synthesis of New 1, 2, 3-triazoles Obtained by 1, 3-Dipolar Cycloaddition Derived from 5-chloro-1-(prop-2-yn-1-yl)indoline-2, 3-dione. *Int.J.Sci. Res.* **2017**,6, 1069-1074.
- [2] Tribak Z.; Ghibate, R.; Skalli M. K.; KandriRodi, Y.; Senhaji, O.; Experimental and Theoretical Study for Corrosion Inhibition in 1M HCl Solution by New 5-Chloroisatin Derivative *Inter. J. Sci. Tech. Eng.* **2016**, 3,257- 262.
- [3] Tribak, Z.;Haoudi, A.; KandriRodi, Y.; Elmsellem, H.; Skalli, M. K.; Ouzidan, Y.; Mazzah, A.; Essassi, E.M.; Synthesis and reactivity of new heterocyclic systems derived from 5-chloro-1H-indole-2,3-dione. *Mor. J. Chem.* **2016**, 4(4), 1157-1163
- [4] Misra, R.N. Rawlins, D.B. Xiao, H. Shan, W. Bursuker, I. Kellar, K.A. Mulheron, J.G. Sack, J.S. Tokarski, J.S. Kimball, S.D. Webster, K.R. 1H-Pyrazolo[3,4-b]pyridine inhibitors of cyclin-dependent kinases, *Bioorg. Med. Chem. Lett.* **2003**, 13, 1133-1136.
- [5] Srivastava A, Pandeya SN: "Indole" a versatile nucleus in the pharmaceutical field. *Int J Curr Pharm Rev Res*, **2011**, 1, 1–17.
- [6] Meijer, L. Flajolet, M. Greengard, P. Pharmacological inhibitors of glycogen synthase kinase 3, *Trends in Pharm. Sci.* **2004**, 25, 471-480.
- [7] Foks, H. Pancechowska-Ksepko, D. Kvdzia, A Zwolska, Z. Janowiec, M. Augustynowicz-Kopec, E. Synthesis and antibacterial activity of 1H-pyrazolo[3,4-b]pyridine and -pyridine derivatives, *Il Farmaco* **60** (2005) 513-517.
- [8] Goda, F.E. Abdel-Aziz, A.A.M. Attef, O.A. Synthesis, antimicrobial activity and conformational analysis of novel substituted pyridines: BF₃-promoted reaction of hydrazine with 2-alkoxy pyridines, *Bioorg. Med. Chem.* **2004**, 12, 1845-1852.
- [9] Kundariya, D.S. Bhesdadia, B.M. Joshi, N.K. Patel, P.K. Synthesis, characterization and pharmacological evaluation of some novel Schiff bases containing 1H-pyrazolo[3,4-b]pyridine moiety, *Int. J. Chem. Tech. Res.* **2011**, 3, 238-243.
- [10] Kini, S.; Swain, S.; Gandhi, A. Synthesis and Evaluation of novel Benzothiazole Derivatives against Human Cervical Cancer cell lines. *Ind. J. Pharm. Sci.* **2007**, 46–50.
- [11] Wang, M.; Gao, M.; Mock, B.; Miller, K; Sledge, G.; Hutchins, G.; Zheng, Q Synthesis of C-11 labelled fluorinated 2-arylbenzothiazoles as novel potential PET cancer imaging agent. *Bioorg. Med. Chem.* **2006**, 14, 8599–8607.
- [12] Pakravan, P., Kashanian S., Khodaei, M.M., Harding, F.J.; Biochemical and pharmacological characterization of isatin and its derivatives: from structure to activity, *Pharmacol. Rep.* **2013**; 65(2), 313-335.
- [13] Sriram, D., Bal, T.R., Yogeewari, P., Synthesis, antiviral and antibacterial activities of isatinmannich bases, *Med. Chem. Res.* **2005**; 14(4), 211- 228.
- [14] Mondal, P., Banerjee, M., Jana, S., Bose, A., Synthesis and evaluation of 1,3 Di-substituted schiff, mannich bases and spiroisatin derivatives, *J. Young Pharm.* **2010**; 2(2), 169-172.
- [15] Wakchaure, N.D., Shejwal, S.S., Deshmukh, V.K, Chaudhari, S.R, Review on Common Methods to Synthesize Substituted 1H-Indole-2, 3-Dione (Isatin) Derivatives and Their Medicinal Significance, *American J. PharmTech Res* **2012**; 2(4): 288-310.
- [16] Malawska, B., New Anticonvulsant Agents, *Curr. Top. Med. Chem.* **2005**; 5(1): 69-85.
- [17] Pandeya, S.N, Sriram, D., Yogeewari, P., Stables, J.P., Anticonvulsant and neurotoxicity evaluation of 5-(un)-substituted isatinimino derivatives, *Pharmazie*, **2001**; 56(11), 875-876
- [18] Tribak, Z.; Haoudi, A.; Skalli, M.K.; KandriRodi, Y.; El Azzouzi, M.; Aouniti, A.; Hammouti, B. Senhaji, O. 5-Chloro-1H-indole-2,3-dione derivative as corrosion inhibitor for mild steel in 1M H₃PO₄: weight loss, electrochemical and SEM studies. *J. Mater. Environ.Sci.* **2017**, 8 (1), 298-309.
- [19] Tribak, Z.;KandriRodi, Y.; Elmsellem, H.; Abdel-Rahman, I.; Haoudi, A.; Skalli, M. K.; Kadmi, Y.; Hammouti, B.; Ali Shariati, M.; Essassi, E. M. 5-chloro-1-octylindoline-2,3-dione as a new corrosion inhibitor for mild steel in hydrochloric acid solution *J. Mater. Environ. Sci.* **2017**, 8 (3) 1116-1127.
- [20] Tribak, Zineb.; KandriRodi, Youssef.; Haoudi, Amal.; Essassi, El Mokhtar.; Capet Frédéric.;

- Zouihri, Hafid 1-Benzyl-5-chloroindoline-2,3-dione. *IUCrData*. 2016, 1(6) 1, x160854.
- [21] Tribak, Zineb.; KandriRodi, Youssef.; Haoudi, Amal.; Essassi, El Mokhtar.; Capet Frédéric.; Zouihri, Hafid, 1-(12-Bromododecyl)-5-chloroindoline-2,3-dione. *IUCrData* 1(6) (2016). 1, x160971.
- [22] Tribak, Zineb.; KandriRodi, Youssef.; Haoudi, Amal.; Essassi, El Mokhtar.; Capet Frédéric.; Zouihri, Hafid, 1-Allyl-5-chloroindoline-2,3-dione. *IUCrData*. 2016, 1(6), 1, x160862.
- [23] Tribak, Zineb.; KandriRodi, Youssef.; Haoudi, Amal.; Essassi, El Mokhtar.; Capet Frédéric.; Zouihri, Hafid, 5-Chloro-1-methylindoline-2,3-dione, *IUCrData*. 2016, 1(6), 1, x160913
- [24] Tribak, Z. El Amin, O. Skalli, M.K. Senhaji, O. Kandri. Rodi, Y. Houssaini. Iraqui, M. N-alkylation methods, Characterization and Evaluation of antibacterial activity of some Novel 5-Chloroisatin Derivatives *Int. J. Eng Res Appl.* 2017, 7(6)21-24
- [25] Tribak, Z.; El Amin, O.; Skalli, M.K.; Senhaji, O.; Kandri. Rodi, Y.; Houssaini. Iraqui, M.; Synthesis, Characterization, and Antibacterial Activity of Some Novel 5-Chloroisatin Derivatives *Int. J. Eng Res Appl.* 2017, 7(6), 66-70.
- [26] Tribak, Z.; KandriRodi, Y.; Haoudi, A.; Skalli, M.K.; Mazzah, A.; Akhazzane, M.; Essassi, E.M.; Cycloaddition 1,3-dipolaire des dérivés de la 5-chloro-1H-indole-2,3 dione: vers de nouvelles isoxazolines et spirodioxazolines *J. Mar. Chim. Heterocycl.* 2016, 16(1), 58-65.
- [27] Tribak, Z. Ghibate, R. Skalli, M.K. KandriRodi, Y. Mrani, D. Aouniti, A. Hammouti, B. Senhaji, O. Synthesis and Characterization of a New Cationic Surfactant Derived from 5-Chloro-1H-indole-2,3-dione In Aqueous Systems, *Int. J. Eng Res Appl.* 2017, 7 (4), 04-08.